

# VU Research Portal

## Winter atmospheric circulation and river discharge in northwest Europe

Bouwer, L.M.; Aerts, J.C.J.H.; Vermaat, J.E.

### **published in**

Geophysical Research Letters  
2006

### **DOI (link to publisher)**

[10.1029/2005GL025548](https://doi.org/10.1029/2005GL025548)

### **document version**

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

### **citation for published version (APA)**

Bouwer, L. M., Aerts, J. C. J. H., & Vermaat, J. E. (2006). Winter atmospheric circulation and river discharge in northwest Europe. *Geophysical Research Letters*, 33 (6)(L06403), 1-4. <https://doi.org/10.1029/2005GL025548>

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

### **E-mail address:**

[vuresearchportal.ub@vu.nl](mailto:vuresearchportal.ub@vu.nl)

# Winter atmospheric circulation and river discharge in northwest Europe

Laurens M. Bouwer,<sup>1</sup> Jan E. Vermaat,<sup>1</sup> and Jeroen C. J. H. Aerts<sup>1</sup>

Received 20 December 2005; revised 2 February 2006; accepted 7 February 2006; published 21 March 2006.

[1] More frequent western atmospheric circulation over Europe results in increased precipitation in winter, and could result in increasing river discharges. We made a quantitative assessment of the impact of variation in atmospheric circulation, defined by the frequency of western circulation in the Großwetterlagen classification system and the North Atlantic Oscillation (NAO) index, on variation in basin-average precipitation and winter discharges (December–February) of eleven large river basins that drain northwest Europe. Annual winter discharges amounts are highly correlated among these rivers (up to  $r^2 = 0.82$ , decreasing with increasing inter-basin distance), which may point to a common atmospheric forcing. The number of days of western atmospheric circulation in winter as indicated by the Großwetterlagen classification system is more closely related ( $r^2 = 0.06$  to  $r^2 = 0.28$ ,  $p < 0.05$  or better) to winter river discharges than the NAO index. We therefore conclude that the frequency of western atmospheric circulation over Europe is a better indicator of climate variability and climate change impacts on river discharges in northwest Europe.

**Citation:** Bouwer, L. M., J. E. Vermaat, and J. C. J. H. Aerts (2006), Winter atmospheric circulation and river discharge in northwest Europe, *Geophys. Res. Lett.*, 33, L06403, doi:10.1029/2005GL025548.


## 1. Introduction

[2] Atmospheric circulation patterns are known to govern variation in precipitation across northwest Europe, particularly in winter [Wibig, 1999]. Some patterns of natural variation in atmospheric circulation have been recognized, of which the North Atlantic Oscillation (NAO) is probably best known. The NAO index is usually defined as the difference of the normalized sea level pressures between the Azores and Iceland, and has been shown to correlate well with precipitation in parts of Europe, most importantly positively in Scandinavia and negatively in the Iberian peninsula [Hurrell, 1995]. Precipitation in locations in western and central Europe that are more distant from the coast, however, is only weakly correlated to the NAO index [Hurrell, 1995; Hurrell and van Loon, 1997; Wibig, 1999].

[3] Since it is widely recognized that precipitation strongly influences river discharges, research has looked

for links between atmospheric circulation patterns and river discharges in Europe. Peterson *et al.* [2002] showed a positive correlation between the NAO index and discharge of northern Eurasian rivers. In a global analysis, Dettinger and Diaz [2000] found significant correlations between the NAO index and river discharge in many locations across Europe. Shorthouse and Arnell [1999], however, showed that strong relationships between the NAO index and winter river discharge in Europe can only be found in Scandinavia and the Iberian Peninsula. Rimbu *et al.* [2004] concluded that a highly smoothed NAO index and annual discharges of the central-eastern located Danube river basin correlate for only 35% of the time for the period 1856–1998.

[4] We made a quantitative assessment of the strength of the relationship between variation in atmospheric circulation and winter river discharges in northwest Europe (December–February). In particular, we address the question to what extent year-to-year variability in winter river discharges can be related to the NAO index. As an alternative we present relationships between river discharge and the frequency of western circulation as recorded in the so-called Großwetterlagen system [Gerstengarbe *et al.*, 1999]. This classification is based on the locations of high and low pressures and ridges and troughs at the 500 hPa level. Although the Großwetterlagen system has been considered a subjective classification scheme, it was found to perform equally well for daily precipitation, when compared to two other objective classification schemes [Buishand and Brandsma, 1997]. Werner *et al.* [2000] showed that the residence time of a particular type of western circulation of the Großwetterlagen system, the West cyclonic circulation, has increased since the 1970s, and some have related this increase to the occurrence of European river floods in the 1990s [Casparly, 1995].



**Figure 1.** Location of the river basins in northwest Europe and the discharge gauging stations (triangles) shown on a 0.5 by 0.5 degree grid. Abbreviations are listed in Table 1.

<sup>1</sup>Institute for Environmental Studies, Faculty of Earth and Life Sciences, Vrije Universiteit, Amsterdam, Netherlands.

**Table 1.** River Discharge Gauging Stations<sup>a</sup>


River	Station	Lat.	Long.	Years	Gauged Area, km <sup>2</sup>	Mean Winter Discharge, m <sup>3</sup> s <sup>-1</sup> (% of Mean Annual Runoff)
Danube (D)	Ceatal Izmail, Romania	45.18°N	28.80°E	1921–1983	807,000	6028 (23%)
Elbe (E)	Neu-Darchau, Germany	53.38°N	11.47°E	1926–2003	131,927	798 (28%)
Garonne (G)	Mas d'Agenais, France	44.42°N	0.24°E	1920–1978 <sup>b</sup>	52,000	912 (37%)
Loire (L)	Montjean, France	47.38°N	0.8°W	1863–1978 <sup>b</sup>	110,000	1352 (40%)
Meuse (M)	Lith, Netherlands	51.81°N	5.45°E	1911–1983	29,000	555 (42%)
Oder (O)	Gozdowice, Poland	52.76°N	14.32°E	1900–1986	109,729	586 (27%)
Rhine (Ri)	Lobith, Netherlands	51.84°N	6.11°E	1901–2001	185,000	2619 (29%)
Rhone (Ro)	Beaucaire, France	43.92°N	4.67°E	1920–1978 <sup>b</sup>	95,590	1919 (28%)
Seine (S)	Paris, France	48.83°N	2.25°E	1928–1978 <sup>b</sup>	44,320	455 (42%)
Weser (We)	Intschede, Germany	52.96°N	9.13°E	1921–1983	37,790	442 (35%)
Wisla (Wi)	Tczew, Poland	54.09°N	18.80°E	1900–1986	194,376	964 (23%)

<sup>a</sup>See also Figure 1.<sup>b</sup>Missing data in 1974 and 1975.

## 2. Methods and Data

[5] Data on monthly river discharges were taken from the RivDIS database at <http://daac.ornl.gov/rivdis/STATIONS.HTM> [Vörösmarty *et al.*, 1998] for nine north-west European rivers (Table 1 and Figure 1). Discharge data for the Elbe were collected from the Arbeitsgemeinschaft für die Reinhaltung der Elbe (ARGE) for the period 1926–2003, and for the Rhine from WaterBase (<http://www.waterbase.nl>) for the period 1901–2001. In addition, winter precipitation data (December–February) for the period 1901–1999 were collected from the CRU TS 2.0 data set, which contains 0.5 by 0.5 degree gridded monthly data based on observations [Mitchell and Jones, 2005]. Monthly precipitation was averaged over each river basin using the basin outlines shown in Figure 1. Basin outlines were taken from Fekete *et al.* [1999]. The NAO index over December–February for the period 1865–2003 was used as developed by Hurrell [1995]. This index is calculated as the difference of normalized sea level pressures between Ponta Delgada in the Azores and Stykkisholmur/Reykjavik in Iceland. The frequency of western circulation (FWC) in winter (December–February) was taken from the Großwetterlagen system for the period 1881–1997 [Gerstengarbe *et al.*, 1999].

[6] First, correlations were made between winter river discharges of different rivers, and a principal component analysis was performed. Secondly, correlations were made between basin-average precipitation and both FWC and the NAO index. Thirdly, regressions were made between winter discharges and precipitation per river basin. Finally,

**Figure 2.** Relationship between coefficients of determination of winter discharges among European rivers and the distance between the geographical centers of the basins.

regressions were made between the two indicators of western atmospheric circulation, the FWC and the NAO index, and annual winter discharges.


## 3. Results

[7] The winter discharges of the river basins in northwest Europe are highly correlated among each other (Table 2). This indicates that drainage systems across northwest Europe behave in a similar manner and may well share a common atmospheric forcing. The correlation is weakest between the rivers located in southwestern Europe (Garonne, Loire) and those in central Europe (Oder, Wisla). Correlation between discharges is probably governed simply by spatial proximity, as correlations are highest among adjacent basins (Figure 2).

[8] Precipitation was found to be important in determining the winter discharge in most river basins (Table 3). Variability in precipitation governed 50% or more of the variation in winter discharge in five out of the eleven basins, and correlations were significant in nine.

[9] Winter values of both the NAO index and FWC show great interannual variation (Figure 3). There is no strong relationship between the NAO index and FWC: the 5-year moving averages of the NAO index and FWC were only weakly correlated, though significantly ( $r^2 = 0.05$ ,  $p < 0.01$ ). Total basin-average winter precipitation was significantly correlated to FWC for nine out of the eleven basins. This basin-average precipitation was not significantly correlated to the NAO index (Table 3).

[10] The winter discharge of nine of the eleven rivers that were studied was significantly correlated to FWC (Table 4). The strongest relationships between FWC and discharge

**Figure 3.** Time series and 5-year moving averages of the NAO index (1865–2003) and the frequency of western atmospheric circulation (FWC, 1881–1997) in winter.

**Table 2.** Coefficients of Determination ( $r^2$ ) of Linear Regressions Among Winter River Discharges<sup>a</sup>

River	Danube	Elbe	Garonne	Loire	Meuse	Oder	Rhine	Rhone	Seine	Weser
Danube	—									
Elbe	0.34 <sup>b</sup>	—								
Garonne	0.18 <sup>b</sup>	0.06	—							
Loire	0.26 <sup>b</sup>	0.13 <sup>c</sup>	0.67 <sup>b</sup>	—						
Meuse	0.23 <sup>b</sup>	0.40 <sup>b</sup>	0.21 <sup>b</sup>	0.44 <sup>b</sup>	—					
Oder	0.30 <sup>b</sup>	0.77 <sup>b</sup>	0.00	0.04 <sup>c</sup>	0.23 <sup>b</sup>	—				
Rhine	0.37 <sup>b</sup>	0.51 <sup>b</sup>	0.28 <sup>b</sup>	0.48 <sup>b</sup>	0.82 <sup>b</sup>	0.31 <sup>b</sup>	—			
Rhone	0.14 <sup>d</sup>	0.09 <sup>c</sup>	0.59 <sup>b</sup>	0.67 <sup>b</sup>	0.28 <sup>b</sup>	0.00	0.45 <sup>b</sup>	—		
Seine	0.18 <sup>d</sup>	0.32 <sup>b</sup>	0.39 <sup>b</sup>	0.75 <sup>b</sup>	0.70 <sup>b</sup>	0.13 <sup>c</sup>	0.73 <sup>b</sup>	0.47 <sup>b</sup>	—	
Weser	0.22 <sup>b</sup>	0.67 <sup>b</sup>	0.05	0.19 <sup>b</sup>	0.70 <sup>b</sup>	0.34 <sup>b</sup>	0.69 <sup>b</sup>	0.12 <sup>b</sup>	0.39 <sup>b</sup>	—
Wisla	0.32 <sup>b</sup>	0.55 <sup>b</sup>	0.01	0.04	0.25 <sup>b</sup>	0.58 <sup>b</sup>	0.32 <sup>b</sup>	0.03	0.08 <sup>c</sup>	0.34 <sup>b</sup>

<sup>a</sup>Length of time series varies according to data availability, see Table 1.<sup>b</sup> $p < 0.001$ .<sup>c</sup> $p < 0.05$ .<sup>d</sup> $p < 0.01$ .

exist for rivers that are strongly correlated to precipitation and drain northwest parts of Europe (Elbe, Meuse, Rhine, Rhone and Weser). Basins that are located more to the south (Garonne, Loire) or to the east (Danube, Oder) have weaker relationships. Year-to-year variability in winter discharge of northwest Europe was not correlated to the winter NAO index (Table 4). Correlations between both winter (December–February) FWC and the NAO index and discharges in December–May do not produce better results, and lagged correlations between both winter (December–February) FWC and the NAO index and discharges in spring (March–May) give even weaker correlations (data not shown here).

[11] A principal component analysis on all rivers for a common period of 1928–1973 shows that the first and second principal components explain 80% of the variance in winter river discharges. In a stepwise multiple linear regression the first principal component was significantly correlated with FWC ( $r^2 = 0.15$ ,  $p < 0.01$ ) but not with the NAO index. The second principal component was again correlated significantly with FWC ( $r^2 = 0.10$ ,  $p < 0.05$ ) and not with the NAO index.

#### 4. Conclusions

[12] Although significant links between the NAO index and river discharges have been found in the south and north

of Europe [Shorthouse and Arnell, 1999; Dettinger and Diaz, 2000; Peterson et al., 2002], our analysis of unsmoothed time series suggests that the correlation between year-to-year winter river discharges and the NAO index in northwest Europe is neither strong, nor significant (Table 4). The frequency of western atmospheric circulation (FWC) as indicated by the Großwetterlagen system appears to be a better indicator for the inter-annual climate variability on river discharges in northwest Europe.

[13] Recent research on North Atlantic climate variability has shown teleconnections with tropical sea surface temperatures [Hurrell et al., 2004]. But smoothed time series of discharges of the Danube River, for instance, are only weakly related to tropical sea surface temperatures [Rimbu et al., 2002, 2004].

[14] The FWC gives a better approximation of variability of river discharges in northwest Europe, as FWC in this system explains 6–28% of annual variation in winter river discharges. Still, this 28% cannot be considered sufficient for prediction purposes and other indicators may need to be included to increase the proportion of explained variance. We conclude that long-term interannual variability in discharge of northwest European rivers

**Table 3.** Coefficients of Determination ( $r^2$ ) of Linear Regressions Between Basin-Average Precipitation and Frequency of Western Atmospheric Circulation (FWC, 97 Years), and the NAO Index (99 Years) in Winter

River	FWC, $r^2$	NAO Index, $r^2$
Danube	0.00	0.00
Elbe	0.27 <sup>a</sup>	0.00
Garonne	0.01	0.00
Loire	0.15 <sup>a</sup>	0.00
Meuse	0.41 <sup>a</sup>	0.00
Oder	0.19 <sup>a</sup>	0.02
Rhine	0.34 <sup>a</sup>	0.00
Rhone	0.08 <sup>b</sup>	0.00
Seine	0.34 <sup>a</sup>	0.00
Weser	0.39 <sup>a</sup>	0.00
Wisla	0.09 <sup>b</sup>	0.00

<sup>a</sup> $p < 0.001$ .<sup>b</sup> $p < 0.01$ .**Table 4.** Coefficients of Determination ( $r^2$ ) of Linear Regressions Between River Discharges and Precipitation, Frequency of Western Atmospheric Circulation (FWC) and the NAO Index in Winter<sup>a</sup>

River	Precipitation		FWC		NAO Index	
	$r^2$	Years	$r^2$	Years	$r^2$	Years
Danube <sup>b</sup>	0.23 <sup>c</sup>	63	0.03	63	0.00	63
Elbe	0.34 <sup>c</sup>	74	0.12 <sup>d</sup>	72	0.00	78
Garonne	0.64 <sup>c</sup>	57	0.09 <sup>c</sup>	57	0.01	57
Loire	0.52 <sup>c</sup>	76	0.06 <sup>c</sup>	96	0.01	112
Meuse	0.60 <sup>c</sup>	73	0.28 <sup>c</sup>	73	0.01	73
Oder	0.31 <sup>c</sup>	86	0.06 <sup>c</sup>	87	0.02	87
Rhine	0.61 <sup>c</sup>	99	0.27 <sup>c</sup>	97	0.00	101
Rhone	0.65 <sup>c</sup>	57	0.17 <sup>c</sup>	57	0.00	57
Seine	0.37 <sup>c</sup>	49	0.07	49	0.00	49
Weser	0.49 <sup>c</sup>	63	0.23 <sup>c</sup>	63	0.02	63
Wisla	0.11 <sup>d</sup>	86	0.17 <sup>c</sup>	87	0.00	87

<sup>a</sup>Length of time series varies according to data availability, see Table 1.<sup>b</sup>21-year averaging for the Danube as performed by Rimbu et al. [2004] leads to coefficients of determination of 0.10,<sup>c</sup> 0.29,<sup>c</sup> and 0.61,<sup>c</sup> respectively.<sup>c</sup> $p < 0.001$ .<sup>d</sup> $p < 0.01$ .<sup>e</sup> $p < 0.05$ .

covaries significantly with North Atlantic climate variability and is much closer linked to the duration of westerly circulation than to large scale north-south pressure differences indicated by the NAO index.

[15] **Acknowledgments.** We thank Michael Bergemann of the Arbeitsgemeinschaft für die Reinhaltung der Elbe for providing us with discharge data of the Elbe. The Climatic Research Unit at the University of East Anglia provided the CRU TS 2.0 data set. Jan Vermaat acknowledges support from the EU FP6 funded ELME project. We thank two anonymous reviewers for their constructive comments. All errors and opinions in this paper are ours.

## References

- Buishand, T. A., and T. Brandsma (1997), Comparison of circulation classification schemes for predicting temperature and precipitation in the Netherlands, *Int. J. Climatol.*, **17**, 875–889.
- Caspary, H. J. (1995), Recent winter floods in Germany caused by changes in the atmospheric circulation across Europe, *Phys. Chem. Earth*, **20**, 459–462.
- Dettinger, M. D., and H. F. Diaz (2000), Global characteristics of stream flow seasonality and variability, *J. Hydrometeorol.*, **1**, 289–310.
- Fekete, B. M., C. J. Vörösmarty, and W. Grabs (1999), Global, composite runoff fields based on observed river discharge and simulated water balances, *Tech. Rep. 22*, World Meteorol. Org., Global Runoff Data Cent., Koblenz, Germany.
- Gerstengarbe, F. W., P. C. Werner, and U. Rüge (1999), *Katalog der Grosswetterlagen Europas (1881–1998) nach Paul Hess und Helmuth Brezowsky*, Potsdam Inst. For Clim. Impact Res., Potsdam, Germany.
- Hurrell, J. W. (1995), Decadal trends in the North Atlantic Oscillation: Regional temperatures and precipitation, *Science*, **269**, 676–679.
- Hurrell, J. W., and H. van Loon (1997), Decadal variations in climate associated with the North Atlantic Oscillation, *Clim. Change*, **36**, 301–326.
- Hurrell, J. W., M. P. Hoerling, A. S. Phillips, and T. Xu (2004), Twentieth century North Atlantic climate change. part I: Assessing determinism, *Clim. Dyn.*, **23**, 371–389.
- Mitchell, T. D., and P. D. Jones (2005), An improved method of constructing a database of monthly climate observations and associated high-resolution grids, *Int. J. Climatol.*, **25**, 693–712.
- Peterson, B. J., R. M. Holmes, J. W. McClelland, C. J. Vörösmarty, R. B. Lammers, A. I. Shiklomanov, I. A. Shiklomanov, and S. Rahmstorf (2002), Increasing river discharge to the Arctic Ocean, *Science*, **298**, 2171–2173.
- Rimbu, N., C. Boroneant, C. Buta, and M. Dima (2002), Decadal variability of the Danube river flow in the lower basin and its relation with the North Atlantic Oscillation, *Int. J. Climatol.*, **22**, 1169–1179.
- Rimbu, N., M. Dima, G. Lohmann, and S. Stefan (2004), Impacts of the North Atlantic Oscillation and the El Niño–Southern Oscillation on Danube river flow variability, *Geophys. Res. Lett.*, **31**, L23203, doi:10.1029/2004GL020559.
- Shorthouse, C. A., and N. W. Arnell (1999), The effects of climatic variability on spatial characteristics of European river flows, *Phys. Chem. Earth, Part B*, **24**, 7–13.
- Vörösmarty, C. J., B. Fekete, and B. A. Tucker (1998), Global river discharge, 1807–1991, version 1.1 (RivDIS dataset), Inst. for the Study of Earth, Oceans, and Space, Univ. of N. H., Durham.
- Werner, P. C., F. W. Gerstengarbe, K. Fraedrich, and H. Oesterle (2000), Recent climate change in the North Atlantic/European sector, *Int. J. Climatol.*, **20**, 463–471.
- Wibig, J. (1999), Precipitation in Europe in relation to circulation patterns at the 500 hPa level, *Int. J. Climatol.*, **19**, 253–269.

J. C. J. H. Aerts, L. M. Bouwer, and J. E. Vermaat, Institute for Environmental Studies, Faculty of Earth and Life Sciences, Vrije Universiteit, De Boelelaan 1087, NL-1081 HV Amsterdam, Netherlands. (laurens.bouwer@ivm.falw.vu.nl)